

Aleuritopteris gongshanensis (Pteridaceae), a New Species from Yunnan, China

Author(s): Gang-Min Zhang, Wen-Li Yang and Xian-Chun Zhang

Source: *Novon*, Vol. 14, No. 4 (Dec., 2004), pp. 513-515

Published by: [Missouri Botanical Garden Press](#)

Stable URL: <http://www.jstor.org/stable/3393554>

Accessed: 06-03-2015 02:27 UTC

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at
<http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Missouri Botanical Garden Press is collaborating with JSTOR to digitize, preserve and extend access to *Novon*.

<http://www.jstor.org>

Aleuritopteris gongshanensis (Pteridaceae), a New Species from Yunnan, China

Gang-Min Zhang

Institute of Botany, Chinese Academy of Sciences, Beijing 100093, People's Republic of China, and Beijing Forestry University, Beijing 100083, People's Republic of China.
gary1967@bjfu.edu.cn

Wen-Li Yang

Hebei Agricultural University, Baoding 071000, People's Republic of China.
yangwl68@sohu.com

Xian-Chun Zhang

Institute of Botany, Chinese Academy of Sciences, Beijing 100093, People's Republic of China. xianchunzhang@yahoo.com.cn

ABSTRACT. *Aleuritopteris gongshanensis* G. M. Zhang, a new species of Pteridaceae from Yunnan Province, China, is described and illustrated. The new species is very similar to *A. pseudoargentea* S. K. Wu, but differs in its ovate-deltate fronds with white farina abaxially, an indusial margin distinctly lacerate, and spores with a reticulate-cristate perine.

Key words: *Aleuritopteris*, China, Pteridaceae.

Aleuritopteris Fée is a middle-sized genus mainly distributed in Asia and Africa (Ching, 1941; Saiki, 1984a, 1984b; Shing & Wu, 1990; Wu, 1981). It is characterized by usually living in a xeric or semi-xeric habitat and having farina abaxially on fronds. This genus was reduced to synonymy of *Cheilanthes* Swartz by some pteridologists (Tryon et al., 1990). Recent studies (Gastony & Rollo, 1995), however, have shown that the species from the Old World are not closely related to those of real *Cheilanthes* from America and should belong to a different group. Here the authors prefer to accept *Aleuritopteris* rather than *Cheilanthes* as the generic name for the group from the Old World.

In revising the Chinese species of *Aleuritopteris*, the authors noticed that some specimens at PYU and PE were erroneously identified as *A. cremea* Ching (= *A. veitchii* (H. Christ) Ching). These specimens are distinguished by having finely dissected fronds, distinctly lacerate indusial margins, and spores with a reticulate-cristate perine. After examining the types of related taxa in *Aleuritopteris*, including *A. veitchii*, *A. pseudoargentea*, and *A. pen-*

tagona Saiki, we confirmed that these specimens were different from the above known species.

***Aleuritopteris gongshanensis* G. M. Zhang, sp. nov.** TYPE: China. Yunnan: Gongshan, Bingzhongluo, under thickets in dry, hot valley, ca. 1650 m, 28 June 1988, W. M. Chu, H. Z. Yan, X. C. Zhang & G. F. Zhang 22789 (holotype, PYU; isotype, PE). Figures 1, 2.

Species *A. pseudoargentea* affinis, sed laminis ovato-deltatis non pentagonis, subtus candido-farinosis, pinnis secundis a basalibus 1.5–4 cm remotis, ambitu conformibus sed minoribus aliquantum, indusiis margine laceratis, perisporis reticulato-cristatis differt.

Rhizomes short, erect, scaly; scales uniformly black or dark brown, concolorous, lanceolate, ca. 0.5 × 6 mm. Leaves monomorphic, clustered, 20–30 cm high; stipes 10–18 cm, longer than laminae, 1–1.5 mm diam., ebeneous, glossy, rounded adaxially, scaly at base (scales similar to those of rhizome); laminae ovate-deltate, caudate at tip, 8–13 × 4.5–8.5 cm, papery or subcoriaceous, adaxially glabrous, abaxially with white farina, tripinnatifid at base; rachises similar to stipes, but sulcate adaxially; pinnae 5 to 8 pairs, usually opposite or sometimes alternate, oblique, sessile, basal pinnae largest, deltate, strongly inequilateral, 4–7 × 4–5 cm, bipinnatifid; pinnules 4 to 7 pairs, alternate; acroscopic pinnules much smaller, usually simple, ca. 8 × 3 mm, obtuse at apex, broadly adnate to costa; proximal basiscopic pinnules greatly enlarged, oblong, to 4 × 1.5 cm, pinnatifid; ultimate segments 4 to 5 pairs, oblong to ovate, ob-

NOVON 14: 513–515. 2004.




Figure 1. *Aleurtopteris gongshanensis* G. M. Zhang. Drawn from the isotype, W. M. Chu et al. 22789 (PE). —a. Habit. —b, c. Rhizome scales. —d. Ultimate segments. —e. Magnification of a segment. —f. Pseudindusium.

tuse at apex, middle ones to 1.5×0.3 cm; second pinnae 1.5–4 cm apart from the basal ones, similar but smaller; upper pairs usually shorter than lower ones. Sori confluent at maturity; pseudindusia narrow, membranous, continuous, with a lacerate mar-

gin. Sporangia 64-spored. Spores subglobose, $45 \times 51 \mu\text{m}$, with a reticulate-cristate perine.

Distribution and habitat. Specimens were collected only in northwestern and central Yunnan.




Figure 2. Spore morphology of *Aleuritopteris gongshanensis*. Spores taken from the isotype, W. M. Chu et al. 22789 (PE). —A. Proximal view, $\times 780$. —B. Distal view, $\times 780$. Bar length = 23.1 μm .

The species usually grows under thickets in dry, hot valleys at ca. 1030–1700 m.

Aleuritopteris gongshanensis is similar to *A. pseudoargentea* and *A. veitchii* in its concolorous rhizome scales, but differs in its ovate-deltate fronds with white farina abaxially, an indusial margin distinctly lacerate, and spores with a reticulate-cristate perine. The new species is also similar to *A. pentagona*, but differs in its concolorous rhizome scales. The main differences between these species are summarized in the following key.

KEY TO *ALEURITOPTERIS GONGSHANENSIS* AND ITS RELATED SPECIES

- 1a. Scales on the rhizome dark brown to black, concolorous.
 - 2a. Indusial margin entire or slightly sinuous; fronds abaxially with yellow or light yellow farina.
 - 3a. Fronds coarsely dissected; spores with a reticulate perine . . . *Aleuritopteris veitchii*
 - 3b. Fronds finely dissected; spores with a cristate perine*Aleuritopteris pseudoargentea*
 - 2b. Indusial margin distinctly lacerate; fronds abaxially with white farina*Aleuritopteris gongshanensis*
- 1b. Scales on the rhizome bicolorous, with dark central stripe and lighter margins*Aleuritopteris pentagona*

Paratypes. CHINA. **Yunnan:** Gongshan, Bingzhongluo, 20 June 1985, W. M. Chu et al. 17681 (PYU), Qinghai-Xizang Exped. 7400 (KUN), 25 May 1960, S. Jiang et al. 8745 (KUN, PE, PYU); nearby Gongshan county

town, 3 Sep. 1988, W. M. Chu et al. 22239 (PYU); Lushui, 14 Oct. 1988, W. M. Chu et al. 22969 (PYU); Yimen, Xiaolüzhi, 20 Oct. 1965, W. M. Chu 4664 (PYU), 26 Apr. 1983, W. M. Chu et al. 15850 (PE, PYU).

Acknowledgments. The first author is grateful to Professor Chu Weiming, curator of the herbarium of Yunnan University, for kind help and hospitality during a visit to PYU. Thanks also to Sun Yingbo for his illustration. This work was supported by a Special Grant for Biological Researches (2001) from the Chinese Academy of Sciences.

Literature Cited

Ching, R. C. 1941. The studies of Chinese ferns—XXXI. Hong Kong Naturalist 10: 194–204.
Gastony, G. J. & D. R. Rollo. 1995. Phylogeny and generic circumscriptions of cheilanthoid ferns (Pteridaceae: Cheilanthesaceae) inferred from *rbcL* nucleotide sequences. Amer. Fern J. 85: 341–360.
Saiki, Y. 1984a. Note on ferns (2). Asiatic species of the *Aleuritopteris farinosa* complex. J. Phytogeogr. Taxon. 32(1): 1–12.
———. 1984b. Note on ferns (3). African and Neotropical species of the *Aleuritopteris farinosa* group. J. Phytogeogr. Taxon. 32(2): 81–90.
Shing, K. H. & S. K. Wu. 1990. Sinopteridaceae. Pp. 97–173 in R. C. Ching & K. H. Shing (editors), Flora Republicae Popularis Sinicae 3(1). Science Press, Beijing.
Tryon, R. M., A. F. Tryon & K. U. Kramer. 1990. *Cheilanthes* Sw. Pp. 242 in K. Kubitzki (editor), The Families and Genera of Vascular Plants, Vol. 1. Pteridophytes and Gymnosperms, K. U. Kramer & P. S. Green (volume editors). Springer-Verlag, Berlin.
Wu, S. K. 1981. A study of the genus *Aleuritopteris* Fée in China. Acta Phytotax. Sin. 19(1): 57–74.