

# *Aphyllodium* (DC.) Gagnep. instead of *Dicerma* DC. (Leguminosae-Papilionoideae: Desmodieae)

Hiroyoshi Ohashi<sup>(1, 2)</sup>

(Manuscript received 7 May 1997; accepted 18 May 1997)

**ABSTRACT:** *Aphyllodium* (DC.) Gagnepain is adopted as the correct generic name instead of *Dicerma* DC. Narrower circumscription of *A. biarticulatum* is adopted, and the following new combinations are made: *Aphyllodium australiense* (Schindl.) H. Ohashi, *A. hispidum* (Schindl.) H. Ohashi, *A. hispidum* var. *collettii* (Schindl.) H. Ohashi, and *A. novoguineense* (Schindl.) H. Ohashi. Key to all the taxa included in the genus is prepared.

**KEY WORDS:** *Aphyllodium*, *Dicerma*, Leguminosae, New combinations.

## INTRODUCTION

Recently *Dicerma* DC. is again replaced by *Aphyllodium* (DC.) Gagnepain (Pedley 1996). This treatment is correct. In the original description, *Dicerma* DC. (1825) includes two sections, i. e., *Phyllodium* (Desv.) DC. and *Aphyllodium* DC. The section *Phyllodium* (Desv.) DC. is, however, published previously as a genus in 1813, hence, according to Art. 52.1 of International Code of Botanical Nomenclature (Greuter *et al.* 1994), the generic name *Dicerma* is superfluous. Nevertheless, the name *Dicerma* was traditionally adopted by Candolle (1826), Wight and Arnott (1834), and Bentham (1852) as the generic name for the section *Aphyllodium* circumscribed by Candolle. This error was already amended by Gagnepain (1916, 1920), but succeeding Schindler (1924), Hutchinson (1964), Ohashi (1973), Ohashi *et al.* (1981), Dy Phon *et al.* (1994), Yang and Huang (1995) and others have used the wrong generic name.

## MATERIALS AND METHOD

Herbarium specimens examined in this study are deposited in A & GH, BM, BO, CAL, K, L, P and TUS. Herbarium acronyms are listed in Holmgren *et al.* (1990).

## RESULTS

In this paper I adopt *Aphyllodium* (DC.) Gagnepain instead of *Dicerma* DC. and change my previous broad specific circumscription of *Dicerma biarticulatum* (L.) DC. in which

1. Biological Institute, Graduate School of Science, Tohoku University, Sendai, 980-77 Japan.

2. Corresponding author.

*Dicerma hispidum* and *Dicerma novoguineense* were included to narrower sense excluding them and *Dicerma biarticulatum* var. *australiense* as distinct.

**Aphyllodium** (DC.) Gagnep. in Notul. Syst. (Paris) **3**: 254. 1916 & in Lecomte, Fl. Gén Indoch. **2**: 609. 1920; Pedley in Rev. Fl. Ceylon **10**: 166. 1996. Type species *Aphyllodium biarticulatum* (L.) Gagnep. (= *Hedysarum biarticulatum* L.).

*Dicerma* sect. *Aphyllodium* DC., Prodr. **2**: 339 (1825) & Mem. Leg. 328 (1826); Wight & Arnott, Prod. Fl. Ind. Orient. 230 (1834). Type species: *Dicerma biarticulatum* (L.) DC. nom. illeg. (= *Hedysarum biarticulatum* L.).

*Desmodium* sect. *Dicerma* (DC.) Benth. in Benth. et Hook. f., Gen. Pl. **1**: 519. 1865 Taubert in Engl. et Pr., Nat. Pflanzenfam. **3**(3): 328. 1894.

*Desmodium* subg. *Dicerma* (DC.) Baker in Hook. f., Fl. Brit. Ind. **2**: 163. 1876.

*Dicerma* DC., Prodr. **2**: 339. 1825 & Mem. Leg. 326. 1826, nom. illeg.; Schindl. in Repert. Spec. Nov. Regni Veg. **20**: 267. 1924; Hutchinson, Gen. Fl. Pl. **1**: 438. 1964; Ohashi in Ginkgoana **1**: 251. 1973; Ohashi et al. in Adv. Leg. Syst. **1**: 299. 1981; Dy Phon et al., Fl. Camb. Laos Vietn. **27**: 43. 1994; Yang et Huang in Fl. Reip. Pop. Sin. **41**: 13. 1995.

### Key to the species

1. Primary bract shorter than the flowers and pods, 2-8 mm long; bracteoles 2-4 mm long; inflorescences lax-flowered; stipules 3-divided at apex, 5-8 mm long; petioles 3-5 mm long; leaflets obovate or elliptic, obtuse at apex, 7-17 mm long, 3-8 mm wide
  2. Primary bracts 3-4 mm long; bracteoles 2-3 mm long; calyx narrowly campanulate, 2.5-3 mm long, tube longer than lobes, the lowest lobe about 1 mm long, almost equal or a little longer than other lobes; appressed pubescent or pilose on inflorescences and pods ..... 2. *A. biarticulatum*
  2. Primary bracts 5-8 mm long; bracteoles ca. 4 mm long; calyx broadly campanulate or funnel-shaped, about 4 mm long, tube shorter than the lobe, the lowest lobe longer than the lateral lobes, about 3 mm long ..... 3. *A. hispidum*
    3. Pods glabrous except for margin; primary bracts 5-6 mm long ..... 3a. var. *collettii*
    3. Pods hirsute; primary bracts 5-8 mm long ..... 3b. var. *hispidum*
1. Primary bract longer than or equal to the flowers and pods, more than 10 mm long; bracteoles 7-9 mm long; inflorescences longer and more densely flowered than above; stipules 3-7-divided at apex, 15-33 mm long; petioles more than 10 mm long; leaflets narrowly elliptic, acute at apex, more than 30 mm long
  2. Primary bracts as long as flowers or pods, 10-13 mm long; bracteoles up to 7 mm long .... 1. *A. australiense*
  2. Primary bracts longer than flowers and pods, 16-17 mm long; bracteoles 7-9 mm long ..... 4. *A. novoguineense*

### 1. *Aphyllodium australiense* (Schindl.) H. Ohashi, *comb. et stat. nov.*

*Dicerma biarticulatum* var. *australiense* Schindl. in Repert. Spec. Nov. Regni Veg. **20**: 267. 1924, p. p., incl. f. *genuinum* Schindl., l. c. 268.

*Desmodium biarticulatum* var. *australiense* (Schindl.) van Meeuwen in Reinwardtia **6**: 247. 1962, p. p.

*Dicerma biarticulatum* subsp. *australiense* (Schindl.) H. Ohashi in Ginkgoana **1**: 255. 1973, p. p.

Distribution: Australia (Queensland, North West Australia).

### 2. *Aphyllodium biarticulatum* (L.) Gagnep., Notul. Syst. (Paris) **3**: 254. 1916 & in Lecomte Fl. Gén. de l'Indoch. **2**: 609. 1920; Pedley in Rev. Fl. Ceylon **10**: 167. 1996.



*Hedysarum biarticulatum* L., Sp. Pl. 747. 1753. Type: Ceylon. *Herb. Hermann* 1: 25 & 3: 15 (BM). *Dicerma biarticulatum* (L.) DC., Prodr. 2: 339. 1825, *nom. illeg.*; Wight & Arnott, Prodr. 230 (1834); Wight, Icon. 2: t. 419. 1843; Benth. in Miq., Pl. Jungh. 217 & 219. 1852; Schindl. in Repert. Spec. Nov. Regni Veg. 20: 267. 1924; Ohashi in Ginkgoana 1: 252. 1973, p. p., excl. subsp. *australiense* (Schindl.) H. Ohashi, p. p. et subsp. *hispidum* (Schindl.) H. Ohashi; Dy Phon et al., Fl. Camb. Laos Vietn. 27: 44. 1994; Ohashi in J. Jpn. Bot. 70: 115. 1995; Yang et Huang in Fl. Reip. Popul. Sin. 41: 13. 1995.

*Desmodium biarticulatum* (L.) Muell., Fragm. Phyt. Austr. 2: 121. 1861; Benth., Fl. Austr. 2: 231. 1864; Baker in Hook. f., Fl. Brit. Ind. 2: 163. 1876; Kurz in J. Asiat. Soc. Beng. 45(2): 232. 1877; Crib, Fl. Siam. Enum. 1: 403. 1928; van Meeuwen in Reinwardtia 6: 246. 1962; Verdc., Man. New Guin. Leg. 392. 1979; Sanjappa, Leg. Ind. 150. 1992.

Distribution: Ceylon, South India, Indo-China (Cambodia, Laos, Thailand, Vietnam), South China (Hainan), Malesia (Java, Celebes, Sumatra), and Australia.

### 3. *Aphyllodium hispidum* (Schindl.) H. Ohashi, *comb. nov.*

Fig. 1



Fig. 1. Holotype of *Dicerma hispidum* Schindl. (CAL).

*Dicerma hispidum* Schindl. in Repert. Spec. Nov. Regni Veg. 20: 269. 1924. Type: Burma. Mengoon. Anderson s. n. on 15 Jan. 1868 (CAL-holotype & isotype).

*D. biarticulatum* subsp. *hispidum* (Schindl.) H. Ohashi in Ginkgoana 1: 257, figs. 69-3 (petals and calyx) & 70-2 (pod and seed). 1973.

### 3a. var. *hispidum*

Distribution: Burma.

### 3b. var. *collettii* (Schindl.) H. Ohashi, *comb. nov.*

Fig. 2

*Dicerma biarticulatum* var. *collettii* Schindl. in Repert. Spec. Nov. Regni Veg. 20: 267. 1924. Type: Burma. Popah District. Collett 881 (CAL-holotype).

Distribution: Burma.




Fig. 2. Holotype of *Dicerma biarticulatum* var. *collettii* Schindl. (CAL)

#### 4. *Aphyllodium novoguineense* (Schindl.) H. Ohashi, *comb. nov.*

*Dicerma novoguineense* Schindl. in Repert. Spec. Nov. Regni Veg. **20**: 268. 1924. Type: New Guinea. Astrolabe Range. *F. H. Brown* 50.

*D. biarticulatum* var. *australiense* f. *longibracteatum* et f. *plumosae* Schindl. in Repert. Spec. Nov. Regni Veg. **20**: 268. 1924.

*Desmodium biarticulatum* var. *australiense* (Schindl.) van Meeuwen in Reinwardtia **6**: 247. 1962, p. p.

*Dicerma biarticulatum* subsp. *australiense* (Schindl.) H. Ohashi in Ginkgoana **1**: 255. 1973, p. p.

Distribution: New Guinea and Australia.

### ACKNOWLEDGEMENT

I am grateful to the director or curators of A & GH, BM, BO, CAL, K, L, and P for their permission to study their collection in their herbaria and/or loan of their specimens.

### LITERATURE CITED

- Baker, J. G. 1876. *Desmodium*. In: Hooker J. D., The Flora of British India **2**: 161-175. Kent.
- Bentham, G. 1852. Leguminosae. In: Miquel F. A. W., Plantae Junghuhnianae. pp. 205-269. Leiden.
- Bentham, G. 1865. *Desmodium*. In: Bentham G. and J. D. Hooker, Genera Plantarum **1**: 519-521. Reeve & Co., London.
- Candolle, A. P. de. 1825. *Dicerma* DC. In: Prodromus Systematics Naturalis Regni Vegetabilis **2**: 339. Paris.
- Candolle, A. P. de. 1826. Revue de la tribu des Hédysarées. In: Mémoires sur la famille des Légumineuses. pp. 295-355. A. Belin, Paris.
- Desvaux, N. A. 1813. *Phyllodium* Desvaux. In: Journal de botanique, appliquee a l'agriculture, a la medecine et aux arts. p. 123. Paris.
- Dy Phon, P., H. Ohashi and J. E. Vidal. 1994. Flore du Cambodge du Laos et du Viêtnam. **27** Leguminosae (Fabaceae), Papilionoideae, Desmodieae. Muséum National d'Histoire Naturelle Laboratoire du Phanérogamie. Paris.
- Gagnepain, F. 1916. Un genre meconnu: *Aphyllodium*. Notul. Syst. (Paris) **3**: 251-255.
- Gagnepain, F. 1920. *Aphyllodium*. In: Lecomte H. (ed.), Flore Générale de l'Indo-Chine **2**: 609-611. Muséum National d'Histoire Naturelle Phanérogamie, Paris.
- Greuter, W. *et al.* 1994. International Code of Botanical Nomenclature (Tokyo Code). Koeltz Sci. Books, Köigstein.
- Holmgren, P. R., N. H. Holmgren and L. C. Barnett. 1990. Index Herbarior. Part I: The Herbaria of the World. 8th edition. International Association for Plant Taxonomy. New York Botanical Garden, New York.
- Hutchinson, J. 1964. Genera of Flowering Plants (Angiospermae) based principally on the Genera Plantarum of G. Bentham and J. D. Hooker. vol. **1**, 516 pp. Oxford Univ. Press London. (Desmodieae pp. 477-484).



- Meeuwen, M. S. Knaap van. 1962. Preliminary revisions of some genera of Malaysian Papilionaceae V - A census of the genus *Desmodium*. *Reinwardtia* 6: 239-276.
- Ohashi, H. 1973. The Asiatic Species of *Desmodium* and Its Allied Genera (Leguminosae). 318 pp., 76 pls. Ginkgoana 1. Academia Scientific Book, Inc., Tokyo.
- Ohashi, H., R. M. Polhill and B. G. Schubert. 1981. Desmodieae. In: Polhill, R. M. and P. H. Raven (eds.), *Advances in Legume Systematics. Part 1*, pp. 292-300. Royal Botanic Gardens, Kew, England.
- Pedley, L. 1996. Fabaceae (Leguminosae) Subfamily Faboideae (Papilionoideae). Tribe Desmodieae. pp. 149-198 (including 11. *Pseudarthria* by V. E. Rudd). Dassanayake, M. D. and W. D. Clayton (eds.), *A Revised Handbook to the Flora of Ceylon* 10. A. A. Balkema, Rotterdam.
- Schindler, A. K. 1924. Über einige kleine Gattungen aus der Verwandtschaft von *Desmodium* Desv. *Repert. Sp. Nov.* 20: 266-286.
- Taubert, P. 1894. Leguminosae. In: Engler A. and K. Prantl (eds.), *Die natürlichen Pflanzenfamilien. III*, 3: 70-396. Leipzig.
- Wight, R. and G. A. W. Arnott. 1834. Leguminosae. In: *Prodromus Florae Peninsula Indiae Orientalis* 1: 178-298. Parbury, Allen, & Co., London.
- Yang, Y.-C. and P.-H. Huang. 1995. Desmodieae. In: Lee S.-K. (ed.), *Fl. Reip. Popul. Sin* 41. Leguminosae (3) Papilionoideae 2. Science Press, Beijing.

## 二節豆屬之學名(豆科—蝶形花亞科：山螞蝗族)

大橋廣好<sup>(1, 2)</sup>

(收稿日期：1997年5月7日；接受日期：1997年5月18日)

### 摘 要

二節豆屬(*Aphyllodium*)分佈在印度、緬甸及澳大利亞北部。大橋廣好原認為 *Dicerma* 與 *Aphyllodium* 有親緣關係。現在根據第一苞片及小苞片間之差異而重新提升亞種為種。二節豆屬學名採用 *Aphyllodium* 而捨棄 *Dicerma*。同時亦採用 *Aphyllodium biarticulatum* 狹義種之觀念，並將亞種提升為種之新組合處理，包括 *A. australiense* (Schindl.) H. Ohashi, *A. hispidum* (Schindl.) H. Ohashi, *A. hispidum* var. *collettii* (Schindl.) H. Ohashi, 及 *A. novoquineense* (Schindl.) H. Ohashi 等。本文亦提供本屬內所有分類群之檢索表，以供參考。

關鍵詞：二節豆屬，豆科，新組合名。

1. 東北大學大學院理學研究科生物學教室，仙台市 980-77，日本。

2. 通訊聯絡員。