

Four new species of *Begonia* L. (Begoniaceae) from Guangxi , China

¹FANG Ding ²WEI Yi-Gang ¹QIN De-Hai

(¹Guangxi Institute of Traditional Medical and Pharmaceutical Sciences , Nanning 530022 , China)

(²Guangxi Institute of Botany , Guangxi Zhuang Autonomous Region and the Chinese Academy of Sciences , Guilin 541006 , China)

Abstract Four new species of the genus *Begonia* L. (Begoniaceae) from Guangxi , China are described and illustrated.

Key words *Begonia* , *B. zhangii* D. Fang & D. H. Qin , *B. jingxiensis* D. Fang & Y. G. Wei , *B. mashanica* D. Fang & D. H. Qin , *B. suboblata* D. Fang & D. H. Qin , new species , Guangxi , China .

In the course of checking specimens of the genus *Begonia* L. (Begoniaceae) from Guangxi , China , the following four species were found to be new to science .

Sect. *Coelocentrum* Irmsch.

1 *Begonia zhangii* D. Fang & D. H. Qin , sp. nov. Fig. 1

张氏秋海棠

Species *B. filiformi* Irmsch. affinis , sed foliis supra annulo lato viridulo-suffuso ornatis , utrinque glandulosis mixtis , petiolis laminis saepe longioribus , scapis brevioribus 11 – 17.5 cm altis , floris masculi feminei tepalis exterioribus omnino majoribus recedit .

Herba perennis , acaulis . Rhizoma repens , villosum , ad nodos radicans , internodiis 4 – 15 mm longis , 4 – 9 mm in diam . Stipulae ovatae vel late ovatae , 9 – 12 mm longae , 5 – 7 mm latae , margine longe ciliatae , dorso secus costam juventute villosae , apice aristatae . Folia 3 – 10 , basalia , petiolis 5 – 22 cm longis crispe brunneo-villosis et glandulosis deinde glabrescentibus ; laminae chartaceae , valde inaequilaterales , ambitu late ovatae , 5.5 – 20 cm longae , 5 – 17 cm latae , apice breviter acuminatae , basi oblique cordatae , margine denticulatae ciliolatae , supra annulo lato viridulo-suffuso ornatae , juventute setulosae et brunneo-glandulosae deinde sensim glabrescentes , subtus dense albido-puncticulatae , brunneo-glandulosae , ad nervos dense crispe brunneo-pubescentes , nervis palmatis 6 – 7 . Scapus ex rhizomate ortus , 11 – 17.5 cm altus , dichasium 2 – 3 (– 6)-ramosum , ramis et pedicellis cum scapo sparse glandulosis et crispe brunneo-pilo-glandulosis , 4 – multi-florum ; bracteae ovato-oblongae vel triangulato-ovatae vel late obovatae , 2.5 – 9 mm longae , 2 – 4 mm latae , margine denticulatae , denticulis apice glanduloso-ciliatis . Tepala rubella . Flores masculi : pedicelli 9 – 13 mm longi ; tepala 4 , exteriora 2 late ovata vel rotunda , 12 – 19 mm longa , 10 – 16 mm lata , extus sparse brunneo-pilo-glandulosa , interiora 2 obovata , 12 – 14 mm longa , 4 – 5 mm lata , glabra , obtusiuscula ; stamina numerosa , basi connata , filamentis 1 mm longis , antheris obovatis , 1.2 – 1.5 mm longis , apice retusis . Flores feminei : pedicelli 7 – 10 mm longi ; tepala 3 , exteriora 2 rotunda , 12 – 15 mm longa , 12 mm lata , extus sparse brunneo-pilo-glandulosa , internum 1 obovatum , 5 mm longum , 2 mm latum , glabrum , subacutum ; stylis 3 , infra medium connati , stigmatibus bifidis contortis papillosis ; ovarium 1-loculatum , placentae segmentis




Fig. 1. *Begonia zhangii* D. Fang & D. H. Qin. A, habit ; B, female flower ; C, capsule ; D, transverse section of capsule. Drawn by X. P. Liao from C. L. Zhang et al. 033 and C. L. Zhang et al. 73310.

2. Capsula pedicello fructifero 14 – 20 mm longo , pendula , anguste ellipsoidea , 15 mm longa , sparse brunneo-pilo-glandulosa , 3-alata , alis lunatis subinaequilongis , ea majore 4.5 mm longa ,

eis duabus minoribus 3 – 4 mm longis. Semina numerosa, minuta. Fl. Mar.

Herb perennial, stemless. Rhizome creeping, villous, rooting at the nodes, internodes 4 – 15 mm long, 4 – 9 mm in diam. Stipules ovate or broadly ovate, 9 – 12 mm long, 5 – 7 mm wide, margin long ciliate, along midrib villous dorsally when young, apex aristate. Leaves 3 – 10, basal; petioles 5 – 22 cm long, crisped brown villous and glandular, then glabrescent; blade papery, very unequally sided, broadly ovate, 5.5 – 20 cm long, 5 – 17 cm wide, apex shortly acuminate, base obliquely cordate, margin denticulate and ciliolate, adaxially adorned with a wide greenish ring, when young setulose and brown glandular, gradually glabrescent, abaxially densely whitish puncticulate and brown glandular, on the nerves densely crisped brown pubescent, palmately 6 – 7-nerved. Scape arising from rhizome, 11 – 17.5 cm high, dichasium 2- or 3-, rarely 6-branched, together with scape sparsely glandulose and crisped brown glandular hairy, 4 – many-flowered; bracts ovate-oblong, triangular-ovate or broadly obovate, 2.5 – 9 mm long, 2 – 4 mm wide, margin denticulate, teeth glandular-ciliate at the tip. Tepals reddish. Male flowers: pedicels 9 – 13 mm long; tepals 4, outer 2 broadly ovate or rounded, 12 – 19 mm long, 10 – 16 mm wide, sparsely brown glandular hairy outside, inner 2 obovate, 12 – 14 mm long, 4 – 5 mm wide, glabrous, somewhat obtuse; stamens numerous; filaments 1 mm long, base connate; anthers obovate, 1.2 – 1.5 mm long, connectives not extended. Female flowers: pedicels 7 – 10 mm long; tepals 3, outer 2 rounded, 12 – 15 mm long, 12 mm wide, sparsely brown glandular hairy outside, inner 1 obovate, 5 mm long, 2 mm wide, glabrous, somewhat acute; styles 3, connate from the middle to bottom; stigmas bifid, contorted, papillate; ovary uniloculed; placentae 3, parietal, bifid. Capsule with stipe 14 – 20 mm long, nodding, narrowly ellipsoid, 15 mm long, sparsely brown glandular hairy, 3-winged, wings crescent-shaped, somewhat unequally long, one larger, 4.5 mm long, the other two smaller, 3 – 4 mm long. Seeds numerous and minute. Flowering in March.

China. Guangxi (广西): Longan (隆安), Dujie (都结), in forests, limestone hill, 1984-03-29, C. L. Zhang et al. (张超良等) 033 (holotype, here designated, GXMI), 042 (GXMI); the same locality, alt. ca. 300 m, 2002-04-04, C. L. Zhang et al. (张超良等) 73310 (frs., GXMI, IBK, PE).

Begonia zhangii is close to *B. filiformis* Irmsch., but differs by having leaf blades adorned with a wide greenish ring above, and mixed glandulose on both surfaces, petioles often longer than leaf blades, scape shorter (11 – 17.5 cm high), and outer tepals of both male and female flowers larger.

The species is named after C. L. Zhang, one of the collectors of the holotype specimen.

Sect. *Diploclinium* (Lindl.) A. DC.

1 *Begonia jingxiensis* D. Fang & Y. G. Wei, sp. nov. Fig. 2

靖西秋海棠

Species nova ob tepala 2 affinis *B. xishuiensi* Ku, a qua rhizomate repenti, foliis 2 – 8, juventute utrinque tomentosis, margine saepe lobatis, lobis late triangulatis 2 – 10 mm longis differt.

Herba perennis, acaulis. Rhizoma repens, noduliforme, saepe ramificatum, juventute villosum, ad nodos radicans, internodiis 3 – 11 mm longis, 4 – 5 mm in diam. Stipulae ovatae, 10 mm longae, 7 mm latae, extus secus costam parce villosae, apice aristatae. Folia 2 – 8, basalia, petiolata; laminae chartaceae, saepe valde inaequilaterales, ambitu late ovatae vel rotundae, 3.5 – 9.5 cm longae, 4 – 11 cm latae, apice breviter acuminatae, basi oblique cordatae, margine denticulatae ciliatae (pilis 1.5 – 5 mm longis) et saepe lobatae, lobis late triangulatis 2 – 10 mm longis,




Fig. 2. *Begonia jingxiensis* D. Fang & Y. G. Wei. A, habit ; B, female flower ; C, capsule ; D, transverse section of capsule. Drawn by X. P. Liao from the type gathering , Y. G. Wei 0215.

utrinque initio brunneo-tomentosae deinde sensim glabrescentes , supra dense foveolatae , subtus ad nervos crispe brunneo-villosae , palmatim 6 – 8-nerves ; petioli robusti 3 – 16 cm longi , 2 – 4 mm in diam. , juventute dense crispe brunneo-villosae et tomentosae deinde subglabri. Scapus saepe ex apice rhizomatis oriundus , 8 – 19 cm altus , angulatus ; dichasium 2 – 3(– 10)-ramosum , ramis cum scapis subglabris , 4 – multi-florum ; bracteae oblongae vel ellipticae , circ. 1 mm longae , 0.7

mm latae, glanduloso-ciliolatae. Tepala rubella. Flores staminati : pedicelli 5 – 15 mm longi ; tepala 2, late ovata vel suborbiculata, 10 – 12 mm longa, 10 – 11 mm lata, extus sparsissime nigro-glandulosa ; stamina numerosa, filamentis 1.5 mm longis, antheris oblongis vel obovatis, 1.2 – 1.5 mm longis, apice retusis. Flores pistillati : pedicelli 5 – 8 mm longi ; tepala 2, oblata, 10 mm longa, 13 mm lata, extus sparsissime nigro-glandulosa ; stylis 3, basi connati, stigmatibus bifidis subcontortis papilloso ; ovarium 3-loculatum, placentis 2-lamellatis. Capsula stipitata, stipite 10 – 20 mm longo, pendula, ovoidea, 11 mm longa, cum alis sparsissime nigro-glandulosa, inaequaliter 3-alata, ala una majore subsemiorbiculata, 8 mm longa, alis ceteris duabus lunatis 3 mm longis. Semina numerosa, minuta. Fl. Oct.

Herb perennial, acaulescent. Rhizome repent, nodule-like, often branched, villous when young, rooting at the nodes, internodes 3 – 11 mm long, 4 – 5 mm in diam. Stipules ovate, 10 mm long, 7 mm wide, along midvein slightly villous dorsally, apex aristate. Leaves 2 – 8, basal, petiolate; blade papery, often very unequally sided, broadly ovate or rounded, 3.5 – 9.5 cm long, 4 – 11 cm wide, apex shortly acuminate, base obliquely cordate, margin denticulate and ciliate (hairs 1.5 – 5 mm long), and often with lobes 2 – 10 mm long and widely triangular, both surfaces first brown tomentose then gradually glabrescent, adaxially densely foveolate, abaxially crisped brown villous on the nerves, palmately 6 – 8-nerved; petioles robust, 3 – 16 cm long, 2 – 4 mm in diam., densely brown crisped villous intermixing with tomentose hairs when young, but subglabrous thereafter. Scape arising from apex of rhizome, 8 – 19 cm high, angular; dichasium 2- or 3-, rarely 10-branched, together with scape subglabrous, 4 – many-flowered; bracts oblong or elliptic, ca. 1 mm long, 0.7 mm wide, glandular-ciliolate. Tepals reddish. Staminate flowers : pedicels 5 – 15 mm long; tepals 2, broadly ovate or suborbicular, 10 – 12 mm long, 10 – 11 mm wide, very scatteredly black glandular outside; stamens numerous; filaments 1.5 mm long; anthers oblong or obovate, 1.2 – 1.5 mm long, connectives not extended. Pistillate flowers : pedicels 5 – 8 mm long; tepals 2, oblate, 10 mm long, 13 mm wide, very scatteredly black glandular outside; styles 3, base connate; stigmas bifid, slightly contorted, papillate; ovary triloculed; placentae axile, bifid. Capsule with stipe 10 – 20 mm long, nodding, ovoid, 11 mm long, together with wings very scatteredly black glandular, unequally 3-winged, the larger one subsemiorbicular, 8 mm long, the other two smaller, crescent-shaped, 3 mm long; seeds numerous, minute. Flowering in October.

China. Guangxi (广西) : Jingxi (靖西), Hurun (湖润), in forests, limestone hill, alt. 450 m, 2002-10-27, Y. G. Wei (韦毅刚) 0215 (holotype, here designated, IBK; isotypes, GXMI, PE).

The species is similar to *B. xishuiensis* Ku in having two tepals, but differs by having rhizome repent, 2 – 8 leaves tomentose on both surfaces when young, leaf blades often lobed, lobes widely triangular, about 2 – 10 mm long. *B. xishuiensis* is glabrous throughout the entire plant and has globose rhizome and only one leaf with lobes 3.8 – 4 cm long.

2 *Begonia mashanica* D. Fang & D. H. Qin, sp. nov. Fig. 3

马山秋海棠

Species nova ob formas foliorum similis *B. leprosae* Hance, sed laminis utrinque fructibusque non scrobiculatis, tepalis 2, fructibus 3-alatis differt.

Herba perennis, acaulis. Rhizoma repens, juventute villosum, ad nodos radicans, internodiis 5 – 10 mm longis, 1.5 – 5 mm in diam. Stipulae latissime ovatae circ. 12 mm in diam., margine fimbriatae, apice aristatae. Folia 2 – 4, basalia, petiolata; laminae chartaceae, aequilaterales interdum subinaequilaterales, reniformes raro ambitu late ovatae, 4 – 7 cm longae, 5 – 11.5 cm




Fig. 3. *Begonia mashanica* D. Fang & D. H. Qin. A , habit ; B , male flower ; C , female flower ; D , transverse section of capsule. Drawn by X. P. Liao from the type gathering , D. Fang & D. H. Qin 31380.

latae , apice rotundae raro subobtusae , basi profunde cordatae interdum subobliquae , margine denticulatae ciliatae (pilis 4 – 5 mm longis) , supra juventute villis brunneolis densis 2.5 – 6 mm longis tectae deinde glabratae , subtus crispe brunneo-villosae , ad nervos densius villosae , palmatim 7-nerves ; petioli 2 – 12 cm longi , crispe brunneo-villosi . Scapus ex apice rhizomatis oriundus , 9.5 – 18 cm altus ; dichasium 2- vel 4-ramosum , ramis et pedicellis cum scapo valde parce atro-brunneo-glandulosis , 4 – multi-florum ; bracteae late obovatae vel oblongae , 2 – 2.5 mm longae , 0.6 – 1.5 mm latae , margine parce atro-brunneo-glandulosae , apice interdum denticulatae , caducae . Tepala rufa . Flores staminati : pedicelli 8 – 13 mm longi , flaccidi ; tepala 2 – 3 , exteriora 2 suborbiculata vel obovatae , 11 – 12 mm longa , 11 – 12.5 mm lata , extus valde parce atro-brunneo-glandulosa , internum 1 anguste obovatum , 8 mm longum , 4 mm latum , acutum ; stamina numerosa ; filamentis 1.5 mm longis ; antheris obovatis , 1 mm longis , apice retusis vel truncatis . Flores pistillati : pedicelli 8 – 10 mm longi , flaccidi ; tepala 2 , late obovata , 13 mm longa , 11 mm lata , extus valde parce atro-brunneo-glandulosa ; stylis 3 , inferne connati , stigmatibus bifidis subcontortis papillosis ; ovarium 3-loculatum , placentis 2-lamellatis . Capsula stipitata , stipite 13 – 15 mm longo , pendula , late ovoidea , 6 mm longa , valde parce atro-brunneo-glandulosa , inaequaliter 3-alata , ala maxima subligulata 6 mm longa , reliquis duabus lunatis 2 – 3 mm longis . Semina numerosa , minuta . Fl. Oct.

Herb perennial , acaulescent . Rhizome repent , villous when young , rooting at the nodes , internodes 5 – 10 mm long , 1.5 – 5 mm in diam . Stipules very broadly ovate , ca. 12 mm in diam . , margin fimbriate , apex aristate . Leaves 2 – 4 , basal , petiolate ; blade papery , equally but sometimes slightly unequally sided , reniform , rarely broadly ovate , 4 – 7 cm long , 5 – 11.5 cm wide , apex rounded , rarely obtuse , base deeply cordate , sometimes slightly oblique , margin denticulate and ciliate (hairs 4 – 5 mm long) , adaxially when young densely covered with brown villous hairs 2.5 – 6 mm long , then glabrate , abaxially crisped brown villous , on the nerves more densely so , palmately 7-nerved ; petioles 2 – 12 cm long , crisped brown villous . Scape arising from apex of rhizome , 9.5 – 18 cm high ; dichasium 2- or 4-branched , together with scape very scatteredly dark brown glandular , 4 – many-flowered ; bracts broadly obovate or oblong , 2 – 2.5 mm long , 0.6 – 1.5 mm wide , margin scatteredly dark brown glandular , apex sometimes denticulate , caducous . Tepals reddish . Staminate flowers : pedicels 8 – 13 mm long , flaccid ; tepals 2 , rarely 3 , outer 2 suborbicular or obovate , 11 – 12 mm long , 11 – 12.5 mm wide , very scatteredly dark brown glandular outside , inner 1 narrowly obovate , 8 mm long , 4 mm wide , acute ; stamens numerous ; filaments 1.5 mm long ; anthers obovate , 1 mm long , connectives not extended . Pistillate flowers : pedicels 8 – 10 mm long , flaccid ; tepals 2 , broadly obovate , 13 mm long , 11 mm wide , very scatteredly dark brown glandular outside ; styles 3 , connate below ; stigmas bifid , slightly contorted , papillate ; ovary triloculed ; placentae axile , bifid . Capsule with stipe 13 – 15 mm long , nodding , broadly ovoid , 6 mm long , very scatteredly dark brown glandular , unequally 3-winged , the larger one subligulate , 6 mm long , the other two smaller , crescent-shaped , 2 – 3 mm long . Seeds numerous and minute . Flowering in October .

China. Guangxi (广西) : Yishan (宜山) , Desheng (德胜) , in forests , 1977-09-29 , Yishan Exped. (宜山队) 4-2-261a (GXMI) ; Mashan (马山) , Yangxu (杨圩) , limestone hill , in cavern , rare , alt. 180 m , 2001-10-24 , D. Fang & D. H. Qin (方鼎 覃德海) 31380 (holotype , here designated , GXMI) .

In the shape of leaves this species is very similar to *B. leprosa* Hance , but differs by having leaves and fruits not scrobiculate , 2 tepals and fruits 3-winged . In *B. leprosa* , there are 4 tepals

and the fruits are clavate and wingless , and the leaves and fruits are obviously scrobiculate when dry.

3 *Begonia suboblata* D. Fang & D. H. Qin , sp. nov. Fig. 4

都安秋海棠

Habitus affinis *B. gulinquigensi* S. H. Huang & Shui , a qua foliis reniformibus juventute supra villis densis 7 – 13 mm longis obtectis , nervis palmatis cum eis lateralibus in vivo albis , floris pistillati tepalis 3 abhorret.

Herba perennis , acaulis. Rhizoma repens , villosum , interdum ramificatum , ad nodos radicans , internodiis 3 – 6 mm longis , 2 – 6 mm diam. Stipulae late vel latissime ovatae , 3 – 5 mm longae , 3 – 6 mm latae , margine fimbriatae , apice aristatae . Folia 2 – 5 , basalia ; petioli 2 – 11 cm longi , juventute dense vetustate parce brunneo-villosi ; laminae chartaceae vel crasse chartaceae , aequilaterales , reniformes , 2 – 4.7 cm longae , 3 – 8 cm latae , basi cordatae vel interdum subobliquae , margine denticulatae et ciliis 1 – 6 mm longis ornatae , supra juventute villis brunneolis densis 7 – 13 mm longis et glandulis atro-brunneolis tectae deinde glabratae , subtus dense albido-puncticulatae et valde parce atro-brunneo-glandulosae , ad nervos brunneo-villosae , nervis palmatis 6 – 8 , cum nervis lateralibus juventute utrinque purpuratis mox supra albentibus . Scapus ex apice rhizomatis oriundus , 5.5 – 14.5 cm altus , cum pedicellis et ramis cymae glaber vel fere glaber ; dichasium 1- vel 2-ramosum , 2 – 4-florum ; bracteae caducae . Tepala intus alba extus rubello-suffusa , glabra . Flores staminati : pedicelli 5 – 15 mm longi ; tepala 4 , 2 exteriora suboblata , 9 – 10 mm longa , 11 – 13 mm lata , 2 interiora obovata , 6 – 7 mm longa , 3 – 3.5 mm lata , obtusiuscula ; stamena numerosa , filamentis 1 – 1.5 mm longis , antheris obovatis , 1.2 mm longis , apice retusis . Flores pistillati : pedicelli 6 – 10 mm longi ; tepala 3 , 2 exteriora suboblata 6 mm longa , 8 mm lata , 1 internum anguste obovatum , 5 mm longum , 2.5 mm latum , obtusiusculum ; styli 3 , basi connati , stigmatibus U-formibus , non contortis , papillosis ; ovarium laevis , 3-loculatum , placentis 2-lamellatis . Capsula pendula , ovoidea , 7 mm longa , glabra , aequaliter 3-alata , alis subsemiorbiculatis 5 mm longis . Semina numerosa , minuta . Fl. Apr.

Herb perennial , acaulescent. Rhizome repent , villous , sometimes branching , rooting at nodes , internodes 3 – 6 mm long , 2 – 6 mm in diam. Stipules broadly or very broadly ovate , 3 – 5 mm long , 3 – 6 mm wide , margin fimbriate , apex aristate. Leaves 2 – 5 , basal ; petioles 2 – 11 cm long , densely villous when young and sparsely villous and brownish when old ; blade papery or thickly papery , equally sided , reniform , 2 – 4.7 cm long , 3 – 8 cm wide , base cordate or sometimes suboblique , margin denticulate and ciliate (hairs 1 – 6 mm long) , adaxially densely brown villous (hairs 7 – 13 mm long) , and dark brown glandular when young , then glabrate , abaxially densely whitish puncticulate and very scatteredly dark brown glandular , brown villous on the nerves , palmately 6 – 8-nerved , palmate nerves together with lateral ones on both surfaces purpurate when young , soon above whitish. Scape arising from apex of rhizome , 5.5 – 14.5 cm high , together with cyme glabrous or almost glabrous ; dichasium 1- or 2-branched , 2 – 4-flowered ; bracts caducous . Tepals white inside , reddish outside , glabrous. Stamine flowers : pedicels 5 – 15 mm long ; tepals 4 , outer 2 suboblata , 9 – 10 mm long , 11 – 13 mm wide , inner 2 obovate , 6 – 7 mm long , 3 – 3.5 mm wide , slightly obtuse ; stamens numerous ; filaments 1 – 1.5 mm long ; anthers obovate , 1.2 mm long , connectives not extended. Pistillate flowers : pedicels 6 – 10 mm long ; tepals 3 , outer 2 suboblata , 6 mm long , 8 mm wide , inner 1 narrowly obovate , 5 mm long , 2.5 mm wide , slightly obtuse ; styles 3 , base connate , stigmas U-shaped , not contorted , papillate ; ovary smooth , triloculed ; placentae axile , bifid. Capsule nodding , ovoid , 7 mm long , glabrous , equally 3-winged , wings subsemiorbicular , 5 mm long. Seeds numerous and minute. Flowering in April.




Fig. 4. *Begonia suboblata* D. Fang & D. H. Qin. A, habit; B, male flower; C, female flower; D, transverse section of capsule. Drawn by X. P. Liao from the type gathering, G. F. Meng 29613.

China. Guangxi (广西): Du'an (都安), Xiaao (下坳), in forests, limestone hill, rare, 1978-04-18, J. Y. Ban (班景益) 4-10-210 (GXMI); the same locality, alt. 760 m, 2002-04-19, G. F. Meng (蒙高飞) 29613 (holotype, here designated, GXMI).

This species resembles *B. gulinquigensis* S. H. Huang & Shui, but differs by having leaf blades reniform, adaxially densely villous with hairs 7–13 mm long when young, both palmate and lateral nerves whitish in the living state, and 3 tepals in pistillate flowers. In *B. gulinquigensis*, the leaf blades are somewhat orbicular, and sparsely setulose on the upper surface, both palmate and lateral veins are green in the living state and its pistillate flowers have 5 tepals.

This species may be cultivated as ornamental plant.

Acknowledgements We thank Professor Wu Zheng-Yi(C. Y. Wu) and Dr. Shui Yu-Min , Kunming Institute of Botany , the Chinese Academy of Sciences , and Dr. Qin Hai-Ning , Institute of Botany , the Chinese Academy of Sciences , for kindly letting us use their literature , and thank Mr. Liao Xin-Pei , Guangxi Institute of Traditional Medical and Pharmaceutical Sciences , for making drawings .

广西秋海棠属(秋海棠科)4 新种

¹ 方 鼎 ² 韦毅刚 ¹ 覃德海

(广西中医药研究所 南宁 530022)

(广西壮族自治区·中国科学院广西植物研究所 桂林 541006)

摘要 描述了广西秋海棠属(秋海棠科)的侧膜胎座组和秋海棠组 4 新种。

关键词 秋海棠属 ; 张氏秋海棠 ; 靖西秋海棠 ; 马山秋海棠 ; 都安秋海棠 ; 新种 ; 中国