

中国椴树科矩圆叶椴一新变种*

A NEW VARIETY OF *TILIA OBLONGIFOLIA* REHD. (TILIACEAE) FROM CHINA

桑植椴 新变种

Tilia oblongifolia Rehd. var. *sangzhiensis* B. R. Liao et W. X. Wang, var. nov.

A var. *oblongifolia* differt ramulis hornotinis et gemmis terminalibus et petiolis fulvo-velutinis et stellato-pilosis, ramulis annotinis leviter fulvo-velutinis et stellato-pilosis, foliis subtus candide stellato-pilosis, in sicco supra atro-viridibus, bracteis utrinque stellato-pilosis.

Hunan(湖南): Sangzhi(桑植), Nature Reserve, Badagong Mountain(八大公山自然保护区), Xiaodingping(小顶坪), in forest, alt. 1600 m, 1994-09-30, B. R. Liao *et al.* (廖博儒等) 1001 (typus, EU; isotypus, in Herb. arium, Yongding Institute of Forest, Zhangjiajie City); Jianzuihe(尖嘴河), in forest, alt. 1450 m, 1995-06-25, B. R. Liao *et al.* 1990; Wanjiatai(万家台), in forest, alt. 1000 m, 1995-09-20, B. R. Liao *et al.* 2001.

本变种与矩圆叶椴主要区别在于嫩枝密被黄褐色茸毛和星状毛,老枝有毛,顶芽有毛,叶下面密被白色星状毛,叶干后上面深绿色,叶柄有毛,苞片两面有星状毛而不同。

王万贤(湖北大学生命科学系,武汉 430062)

Wang Wan-xian (Department of Life Science, Hubei University, Wuhan, 430062)

廖博儒(张家界市永定区林业科学研究所,张家界 416600)

Liao Bo-ru (Yongding Institute of Forest, Zhangjiajie City, Zhangjiajie, 416600)