

窄叶山血丹(紫金牛科)一新变种

马晓洁¹, 胡启明², 冯志坚¹

(1. 华南农业大学林学院, 广州 510640; 2. 中国科学院华南植物园, 广州 510650)

摘要: 报道了紫金牛科紫金牛属的一个新变种 窄叶山血丹 *Ardisia lindleyana* D. Dietr. var. *angustifolia* C. M. Hu & X. J. Ma, 并比较了此变种与本属中具有类似叶形种类的区别。

关键词: 紫金牛属 窄叶山血丹 紫金牛科 新变种 广东

中图分类号: Q949.773.1

文献标识码: A

文章编号: 1005-3395(2007)06-0551-02

A New Variety of *Ardisia lindleyana* D. Dietr. (Myrsinaceae) from Guangdong, China

MA Xiao-jie¹, HU Qi-ming², FENG Zhi-jian¹

(1. College of Forestry, South China Agricultural University, Guangzhou 510640, China;

2. South China Botanical Garden, the Chinese Academy of Science, Guangzhou 510650, China)

Abstract: A new variety, *Ardisia lindleyana* var. *angustifolia* C. M. Hu & X. J. Ma is described. It differs from the type variety by its leaves being linear-lanceolate, 7-13.5 cm long, only 4-7 mm broad. Superficially, it strikingly resembles *A. ensifolia* E. Walker and *A. quinqueгона* Blume var. *salicifolia* (E. Walker) C. M. Hu & J. E. Vidal, but can be easily distinguished by its rusty puberulent branchlets, inflorescences terminal on leafy branchlets, and leaves with marginal glands.

Key words: *Ardisia Swartz*; *Ardisia lindleyana* var. *angustifolia*; Myrsinaceae; New variety; Guangdong

Since the publication of the Myrsinaceae treatments in the *Flora Reipublicae Popularis Sinicae*^[1] and *Flora of China*^[2], the authors carried out a project of investigating and evaluating the resources of *Ardisia* species in Guangdong Province, China. In the process, a plant of the genus with very narrow linear-lanceolate leaves was found (Fig. 1). Although it strikingly resembles *A. ensifolia* E. Walker and *A. quinqueгона* Blume var. *salicifolia* (E. Walker) C. M. Hu & J. E. Vidal in aspect, its affinity is certainly with *A. lindleyana* D. Dietr.. Thus it is described as a new variety of the species.

Ardisia lindleyana D. Dietr. var. *angustifolia* C. M. Hu & X. J. Ma, var. nov.

A var. *lindleyanae* recedit foliis lineari-lanceolatis, 7-13.5 cm longis et 4-7 mm latis, margine valde revolutis. Primo visu *A. ensifoliae* E. Walker et *A. quinqueгонаe* var. *salicifoliae* (E. Walker) C. M. Hu & J. E. Vidal maxime similis, tamen non arcte affinis, a quibus ramulis ferrugineo-puberulis, foliis glandulis marginalibus instructis, inflorescentiis in ramulis foliatis terminatis differt.

Guangdong(广东): Enping(恩平), Qingwan(清湾), Qixingkeng(七星坑), in thin forests, alt. 600 m, 1998-11-28, B. H. Chen(陈炳辉) 1885 (Holotype, IBSC); the same locality, on upper hillsides and mountain ridges, alt. 600 m, 1995-12-10, Z. M. Wu(吴志敏) S426 (CANT).



Fig. 1 窄叶山血丹果枝 Fruit branches of *Ardisia lindleyana*
D. Dietr. var. angustifolia C. M. Hu & X. J. Ma

References

- [1] Chen C. Flora Reipublicae Popularis Sinicae Tomus 58 [M]. Beijing Science Press, 1979:35- 98.(in Chinese)
- [2] Chen C. Myrsinaceae [M]// Wu Z Y, Raven P H. Flora of China, Vol. 15. Beijing: Science Press, St. Louis: Missouri Botanical Garden Press, 1996:10- 29.